

RUNNING HEAD: BIG FIVE GWAS

Robust inference and widespread genetic correlates from a large-scale genetic association study of human personality

Ted Schwaba^{1†}, Margaret L. Clapp Sullivan², Wonuola A. Akingbuwa^{3,4}, Kerli Ilves⁵, Peter T. Tanksley⁶, Camille M. Williams⁷, Yavor Dragostinov^{2,8}, Wangjingyi Liao⁹, Lindsay S. Ackerman¹, Josephine C. M. Fealy¹, Gibran Hemani^{10,11}, Javier de la Fuente², Priya Gupta¹², Murray B. Stein¹³, Joel Gelemler¹², Daniel F. Levey¹², Urmo Vösa⁵, Liisi Ausmees¹⁴, Anu Realo^{14,15,16}, Estonian Biobank Research Team⁵, Mariliis Vaht⁵, Jüri Allik¹⁴, Tõnu Esko⁵, René Mõttus^{14,17}, Uku Vainik^{5,14,18}, Gudrun A. Jonsdottir¹⁹, Gudmar Thorleifsson¹⁹, Árni Freyr Gunnarsson¹⁹, Gyda Bjornsdottir¹⁹, Thorgeir E. Thorgeirsson¹⁹, Hreinn Stefansson¹⁹, Kari Stefansson¹⁹, Rosa Cheesman^{20,21}, Qi Qin²⁰, Elizabeth C. Corfield^{10,21,22}, Helga Ask^{20,21}, Fartein Ask Torvik^{20,23}, Eivind Ystrom^{20,21,24}, Martin Tesli^{25,26}, Dorret I. Boomsma²⁷, Eco J. C. de Geus^{3,4}, Jouke-Jan Hottenga³, Dener Cardoso Melo²⁸, Harold Snieder²⁹, Catharina A. Hartman^{17,28}, Charley Xia^{17,30}, Archie Campbell³¹, Michelle Luciano¹⁷, Ian J. Deary^{17,30}, W. David Hill^{17,30}, Seon-Kyeong Jang³², Scott I. Vrieze^{33,34}, Gonçalo Abecasis³⁵, Michelle K. Lupton^{36,37,38}, Brittany L. Mitchell^{36,37,38}, Petra V. Viher³⁶, Lucía Colodro-Conde³⁶, Nicholas G. Martin³⁶, Sarah E. Medland³⁶, Eske M. Derks³⁶, Briar Wormington³⁶, Jaakko Kaprio³⁹, Karri Silventoinen⁴⁰, Teemu Palviainen³⁹, Agnieszka Gidziela⁹, Kaili Rimfeld^{41,42}, Robert Plomin⁴², Margherita Malanchini⁹, Danielle M. Dick^{43,44}, Fazil Aliev⁴⁴, COGA Collaborators, The Spit for Science Working Group⁴⁶, Laura W. Wesseldijk^{45,46,47}, Fredrik Ullén^{45,46}, Miriam A. Mosing^{46,48,49}, Henry R. Kranzler⁵⁰, Yaira Nunez⁵¹, Sarah Beck⁵¹, Renato Polimanti⁵¹, Tobias Edwards³³, Alexandros Giannelis³³, Emily A. Willoughby³³, James J. Lee³³, Matt McGue³³, Antonio Terracciano^{52,53}, Michele Marongiu⁵⁴, Edoardo Fiorillo⁵⁴, Francesco Cucca⁵⁵, Angelina R. Sutin⁵⁶, Peter J. van der Most²⁹, Albertine J. Oldehinkel²⁸, Tina Kretschmer⁵⁷, Andrey A. Shabalin⁵⁸, Anna R. Docherty⁵⁸, Robert F. Krueger³³, Colin D. Freilich³³, Binisha H. Mishra^{59,60,61}, Terho Lehtimäki^{59,60,61}, Olli T. Raitakari^{62,63,64,65}, Mika Kähönen^{60,66}, Aino Saarinen⁶⁷, Henrik Dobewall⁶⁸, Liisa Keltikangas-Järvinen⁶⁷, Klaus Berger⁶⁹, Marisol Herrera-Rivero^{69,70}, Fabian Streit^{71,72,73}, Swapnil Awasthi⁷⁴, Stephanie H. Witt⁷⁵, Johanna Tuhkanen⁷⁶, Katri Räikkönen⁷⁶, Johan G. Eriksson^{77,78,79}, Jari Lahti^{76,78}, Gail Davies³⁰, Paul Redmond³⁰, Adele Taylor³⁰, Janie Corley³⁰, Tom C. Russ^{80,81}, Marina Ciullo⁸², Teresa Nutile⁸², Jun Ding⁸³, Yong Qian⁸³, Toshiko Tanaka⁸³, Luigi Ferrucci⁸³, Lea Zillich⁷⁵, Lea Sirignano⁷⁵, K. Paige Harden², Erhan Genç⁸⁴, Patrick D. Gajewski⁸⁵, Stephan Getzmann⁸⁵, Christoph Fraenz⁸⁴, Javier E. Schneider Peñate⁸⁶, Stefanie Lis^{87,88}, Alisha S. M. Hall^{75,89,90}, Christian Schmahl⁸⁷, Sabine C. Herpertz⁹¹, Abdel Abdellaoui^{47,92}, Michel G. Nivard^{10,11*}, Elliot M. Tucker-Drob^{2,8*}

1. Department of Psychology, Michigan State University, East Lansing, MI, USA
2. Department of Psychology, University of Texas at Austin, Austin, TX, USA
3. Department of Biological Psychology, Faculty of Behavioural and Movement Sciences, Vrije Universiteit, Amsterdam, the Netherlands.
4. Amsterdam Public Health Research Institute, Amsterdam University Medical Centre, Amsterdam, the Netherlands.
5. Institute of Genomics, University of Tartu, Estonia
6. ALERRT Center, Texas State University
7. Laboratoire de Sciences Cognitives et Psycholinguistique (ENS, EHESS, CNRS), École Normale Supérieure, PSL University, Paris, France
8. Population Research Center, University of Texas at Austin, Austin, TX, USA
9. School of Biological and Behavioral Sciences, Queen Mary University of London, London, UK
10. Medical Research Council Integrative Epidemiology Unit, University of Bristol, Bristol, UK
11. Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK
12. Yale University, Veterans Affairs Connecticut Healthcare System
13. University of California San Diego, Veterans Affairs San Diego Healthcare System
14. Institute of Psychology, University of Tartu, Estonia
15. Department of Psychology, University of Warwick, UK
16. School of Educational Sciences, Tallinn University, Estonia
17. Department of Psychology, University of Edinburgh, Edinburgh, UK
18. Montreal Neurological Institute, McGill University, Canada
19. deCODE genetics/Amgen, Reykjavik, Iceland
20. PROMENTA Research Center, Department of Psychology, University of Oslo, Oslo, Norway
21. PsychGen Centre for Genetic Epidemiology and Mental Health, Division of Public Health and Prevention, Norwegian Institute of Public Health, Oslo, Norway
22. Psychiatric Genetic Epidemiology Group, Research Department, Lovisenberg Diaconal Hospital, Oslo, Norway
23. Centre for Fertility and Health, Norwegian Institute of Public Health, Oslo, Norway
24. Centre for Research on Equality in Education, Faculty of Educational Sciences, University of Oslo, Oslo, Norway
25. Department of Mental Health, Norwegian Institute of Public Health, Oslo, Norway

26. Department of Psychiatry, Østfold Hospital, Norway
27. Department of Complex Trait Genetics, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands
28. Interdisciplinary Center Psychopathology and Emotion Regulation (ICPE), University of Groningen, University Medical Center Groningen, Groningen, The Netherlands
29. Epidemiology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands
30. Lothian Birth Cohort Studies, Department of Psychology, University of Edinburgh, Edinburgh, UK
31. Centre for Genomic and Experimental Medicine, Institute of Genetics and Cancer, University of Edinburgh, Edinburgh, UK
32. Department of Computational Medicine, University of California Los Angeles, Los Angeles
33. Department of Psychology, University of Minnesota
34. Minnesota Center for Twin and Family Research
35. Regeneron Genetics Center, Regeneron Pharmaceuticals
36. QIMR Berghofer Medical Research Institute, Brisbane, QLD, Australia
37. Faculty of Medicine, The University of Queensland, Brisbane, QLD, Australia
38. School of Biomedical Sciences, Queensland University of Technology, Brisbane, QLD, Australia.
39. Institute for Molecular Medicine FIMM, University of Helsinki, Helsinki, Finland
40. Population Research Unit, Faculty of Social Sciences, University of Helsinki, Helsinki, Finland
41. Department of Psychology, Royal Holloway University of London, Egham, UK
42. Social, Genetic, & Developmental Psychiatry Centre, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, UK
43. Rutgers Addiction Research Center, Brain Health Institute, Rutgers University
44. Department of Psychiatry, Rutgers Robert Wood Johnson Medical School
45. Department of Neuroscience, Karolinska Institutet, Stockholm, Sweden
46. Department of Cognitive Neuropsychology, Max Planck Institute for Empirical Aesthetics, Frankfurt am Main, Germany
47. Department of Psychiatry, Amsterdam University Medical Centre, University of Amsterdam
48. Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden
49. Melbourne School of Psychological Sciences, University of Melbourne, Australia
50. Department of Psychiatry, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA, United States
51. Department of Psychiatry, Yale School of Medicine, New Haven CT, United States
52. Department of Geriatrics, College of Medicine, Florida State University, Tallahassee
53. Laboratory of Behavioral Neuroscience, National Institute on Aging, Baltimore
54. Institute for Genetic and Biomedical Research, Consiglio Nazionale delle Ricerche (CNR), Cagliari, Italy
55. Department of Biomedical Sciences, University Study of Sassari, Sassari, Italy
56. Department of Behavioral Sciences and Social Medicine, College of Medicine, Florida State University, Tallahassee
57. Pedagogical Sciences, Faculty of Behavioural and Social Sciences, University of Groningen, Groningen, The Netherlands
58. Huntsman Mental Health Institute, Department of Psychiatry, University of Utah
59. Department of Clinical Chemistry, Faculty of Medicine and Health Technology, Tampere University, Tampere, Finland
60. Finnish Cardiovascular Research Center Tampere, Faculty of Medicine and Health Technology, Tampere University, Tampere, Finland
61. Department of Clinical Chemistry, Fimlab Laboratories, Tampere, Finland
62. Department of Clinical Physiology and Nuclear Medicine, Turku University Hospital, Turku, Finland
63. Centre for Population Health Research, University of Turku and Turku University Hospital, Turku, Finland
64. Research Centre of Applied and Preventive Cardiovascular Medicine, University of Turku, Turku, Finland
65. InFLAMES Research Flagship, University of Turku, Turku, Finland
66. Department of Clinical Physiology, Tampere University Hospital, Tampere Finland
67. Faculty of Medicine, Department of Psychology and Logopedics, University of Helsinki, Helsinki, Finland
68. Finnish Institute for Health and Welfare
69. Institute of Epidemiology and Social Medicine, University of Münster, Germany
70. Department of Genetic Epidemiology, Institute of Human Genetics, Germany
71. Hector Institute for Artificial Intelligence in Psychiatry, Central Institute of Mental Health, Medical Faculty Mannheim, Heidelberg University, Mannheim, Germany
72. Department of Psychiatry and Psychotherapy, Central Institute of Mental Health, Medical Faculty Mannheim, Heidelberg University, Mannheim, Germany
73. German Center for Mental Health (DZPG), Partner Site Mannheim - Heidelberg - Ulm, Germany
74. Department of Psychiatry and Psychotherapy, Campus Mitte Berlin, Charité Universitätsmedizin Berlin, Berlin, Germany
75. Department of Genetic Epidemiology in Psychiatry, Medical Faculty Mannheim/Heidelberg University, Central Institute of Mental Health, Mannheim, Germany
76. Department of Psychology, University of Helsinki, Helsinki, Finland
77. Department of General Practice and Primary Health Care, Tukholmankatu 8 B, University of Helsinki, Finland
78. Folkhälsan Research Centre, Helsinki, Finland
79. National University Singapore, Yong Loo Lin School of Medicine, Department of Obstetrics and Gynecology, Singapore, Singapore
80. Alzheimer Scotland Dementia Research Centre, University of Edinburgh, United Kingdom

81. Division of Psychiatry, Centre for Clinical Brain Sciences, University of Edinburgh, United Kingdom
82. Institute of Genetics and Biophysics "Adriano Buzzati-Traverso" - CNR, Naples, Italy
83. Translational Gerontology Branch, National Institute on Aging, Baltimore
84. Department of Psychology and Neurosciences, Neuroimaging and Interindividual Differences, Leibniz Research Centre for Working Environment and Human Factors (IfADo), Dortmund, Germany
85. Department of Ergonomics, Networking Group Aging, Leibniz Research Centre for Working Environment and Human Factors (IfADo), Dortmund, Germany
86. Department of Psychology, Neuropsychology, Ruhr University Bochum, Bochum, Germany
87. Department of Psychosomatic Medicine and Psychotherapy, Medical Faculty Mannheim/Heidelberg University, Central Institute of Mental Health, Mannheim, Germany
88. Department of Clinical Psychology, Medical Faculty Mannheim/Heidelberg University, Central Institute of Mental Health, Mannheim, Germany
89. Department of Affective Disorders, Aarhus University Hospital, Aarhus, Denmark
90. Department of Clinical Medicine, Aarhus University, Aarhus, Denmark
91. Department of Psychosomatic General Psychiatry, Center for Psychosocial Medicine, Medical Faculty Mannheim/Heidelberg University, Heidelberg, Germany
92. Amsterdam Neuroscience, Amsterdam, The Netherlands

* M.G.N. and E.M.T-D. jointly supervised this work.

† Send correspondence regarding this article to tedschwaba@gmail.com

Abstract

Personality traits describe stable differences in how individuals think, feel, and behave and how they interact with and experience their social and physical environments. We assemble data from 46 cohorts including 611K-1.14M participants with European-like and African-like genomes for genome-wide association studies (GWAS) of the Big Five personality traits (extraversion, agreeableness, conscientiousness, neuroticism, and openness to experience), and data from 51K participants for within-family GWAS. We identify 1,257 lead genetic variants associated with personality, including 823 novel variants. Common genetic variants explain 4.8%-9.3% of the variance in each trait, and 10.5%-16.2% accounting for measurement unreliability. Genetic effects on personality are highly consistent across geography, reporter (self vs. close other), age group, and measurement instrument, and we find minimal spousal assortment for personality in recent history. In stark contrast to many other social and behavioral traits, within-family GWAS and polygenic index analyses indicate little to no shared environmental confounding in genetic associations with personality. Polygenic prediction, genetic correlation, and Mendelian randomization analyses indicate that personality genetics have widespread, potentially causal associations with a wide range of consequential behaviors and life outcomes. The genetic architecture of personality is robust and fundamental to being a human.

Introduction

Personality is defined by relatively stable patterns of thinking, feeling, and behaving that vary across individuals (Roberts and Yoon, 2022). Decades of research have demonstrated that human personality can be meaningfully and efficiently summarized by five broad trait factors known as the *Big Five*: Extraversion (which encompasses traits such as sociability, assertiveness, and energy level), agreeableness (compassion, respectfulness, and trust), conscientiousness (organization, productiveness, and responsibility), neuroticism (anxiety, depression, and volatility) and openness to experience (imagination, curiosity, and aesthetic sensitivity) (John et al., 2008; Matthews et al., 2009; Soto & John, 2017). The Big Five predict academic performance and educational attainment, labor market outcomes, and career choices, at comparable levels to cognitive ability and socioeconomic status (Anni et al., 2024; Borghans et al., 2016; Bucher et al., 2019; Deary et al., 2010; Hampson, 2012; Roberts et al., 2007; Soto, 2019). They relate to a wide range of health behaviors, such as eating habits, drug use, and physical activity (Bogg & Roberts, 2004; De Moor et al., 2006; Hampson et al., 2007; Willroth et al., 2023) and to disease burden (Yoneda et al., 2023) and longevity (Beck & Jackson, 2022; Graham et al., 2017). The Big Five are so closely linked to mental health that modern taxonomies of psychopathology are foundationally informed by patterns of personality trait variation (Kotov et al., 2010; Widiger et al., 2019). And they predict relationships and social behavior (Antonoplis & John, 2017; Landis, 2016), residential mobility (Jokela, 2020; Jokela et al., 2015), and political preferences (Duckitt & Sibley, 2010). Given the fundamental relevance of personality to core phenomena studied across scientific disciplines, understanding the molecular genetics of personality is crucially important for developing biopsychosocial models of both human behavior and life outcomes.

Here we report results from the Revived Genomics of Personality Consortium (ReGPC), a collaboration across 46 cohorts incorporating between 611K and 1.14M participants with European-like (EUR) and African-like (AFR) genomes (National Academies of Sciences, Engineering, and Medicine, 2023) per Big Five trait. This effort represents a substantial boost in statistical power for genetic discovery over past personality GWAS efforts, increasing the number of significant loci relative to most recent work (Gupta et al., 2024) from 3 to 131 for conscientiousness, 4 to 39 for agreeableness, 8 to 126 for openness to experience, 14 to 258 for extraversion, and 224 to 703 for neuroticism. Though personality traits often vary on average across groupings of people (Bleidorn et al., 2022; Ebert et al., 2022), we find that the genetic etiology of each trait is highly consistent across geography, reporter perspective (self vs. close other), age, and measurement instrument. We empirically confirm the strong relationship between a measurement's internal consistency and its heritability; after accounting for measurement error, heritability estimates for the Big Five are substantially greater. We report results of within-family GWAS of the Big Five in over 50K participants, which allow us to parse genetic effects from environmental confounds. We find that the genetic architecture of the Big Five is only minimally confounded by population stratification, dynastic, and assortative mating effects that commonly bias GWAS of other behavioral traits (Howe et al., 2022; Tan et al., 2024; Young et al., 2022). Polygenic indices (PGIs) constructed from each primary GWAS robustly predict personality in five independent cohorts, with no identifiable reduction in predictive accuracy when compared within families. Across the Big Five, common variant heritability estimates are largely indistinguishable when estimated using within-family vs. population GWAS, and population-level and within-family genetic effects are highly genetically correlated

(mean $r_g = .86$). We characterize the functional genomics of each of the Big Five and provide evidence of negative selection on the genetics of personality. Finally, we extensively characterize personality's wide-ranging genetic correlates and consequences across social relevant behaviors and life outcomes ranging from psychopathology and physical health to research participation and residential mobility, highlighting a central role of personality genetics in understanding the human experience.

Results

Population-level Association Meta-Analysis

We performed population-level GWAS meta-analysis of each Big Five trait in genetic-similarity-stratified and genetic-similarity-combined cohorts of EUR ($k = 46$) and AFR ($k = 10$) participants. In each cohort analysts conducted GWAS of each available Big Five measurement instrument using a standard protocol across the 22 autosomes and, when available, the X-chromosome (**Supplementary Tables S1-S3**). We jointly analyzed results using inverse-variance weighted meta-analysis, resulting in pooled associations for ~ 10 million Single Nucleotide Polymorphisms (SNPs) for each Big Five trait.

We identified 1,257 approximately independent ($r^2 < .10$ within a 250kb range) genome-wide significant (GWS; $p < 5 \times 10^{-8}$) SNPs associated with the Big Five traits, an increase of 3 to 43 times over the most recent findings for each trait (**Table 1; Supplementary Tables S4-S11**). Of these lead SNPs, 823 (65%) were found in novel loci not previously associated with variation in the respective trait. Manhattan plots of these results are displayed in **Figure 1 A-E**. Demonstrating the independence between the Big Five, 82% of genomic loci containing a GWS SNP were associated with only a single trait (**Supplementary Table S12**), and the average absolute genetic correlation among Big Five traits among EUR participants was .19 (**Figure 1 F**), which is similar to estimates of phenotypic correlations (Soto & John, 2017).

Table 1. Summary of Population-Level Genome-Wide Association Study Results

Trait	N _{Total} (N _{AFR})	Mean χ^2	Previous Lead SNPs	This Study Lead SNPs	EUR SNP Heritability (h^2_{SNP})				Mean LDSC genetic correlation across grouped cohorts				
					LDSC Intercept (EUR/AFR)	h^2_{SNP} estimate from GWAS meta-analysis	RE-Meta of individual h^2_{SNP} estimates	RE-Meta disattenuated for measurement ins. error	Geographic region	Measure- ment ins.	Reporter perspective	Military service	Age group
Extraversion	662,617 (43,201)	1.59	17	258	1.00/1.01	9.3% (0.3)	10.6% (1.3)	16.2% (1.0)	.90	.93	.92	.84	1.00
Agreeableness	611,037 (43,482)	1.29	4	39	1.02/1.00	4.8% (0.2)	7.4% (1.0)	10.5% (1.0)	.65	.68	.95	.73	.70
Conscientiousness	641,167 (43,120)	1.45	3	131	1.03/1.02	7.1% (0.2)	7.9% (1.0)	11.0% (1.0)	.86	.90	.68	.80	.79
Neuroticism	1,136,711 (48,110)	2.02	224	703	0.99/1.02	8.3% (0.3)	8.4% (0.8)	12.0% (0.8)	.99	.97	.77	.84	.74
Openness to experience	611,985 (43,498)	1.43	8	126	1.03/1.01	7.4% (0.3)	9.4% (1.6)	14.5% (1.2)	.92	.82	.87	.78	.88

Note. h^2_{SNP} = Single Nucleotide Polymorphism (SNP) heritability. EUR = Individuals with European-like genomes. AFR = Individuals with African-like genomes. RE-meta = random-effects meta-analysis of cohort-specific h^2_{SNP} estimates. LDSC = Linkage Disequilibrium Score regression. Measurement ins. = Measurement instrument. Lead SNPs = number of genome-wide significant SNPs ($p < 5 \times 10^{-8}$) that are independent (Linkage Disequilibrium $r^2 < .10$) of other lead SNPs. Prev. = lead SNPs identified in Gupta and colleagues (2024) across all analyses. h^2_{SNP} and genetic correlations were estimated using data from EUR samples using LDSC. Numbers in parentheses represent standard errors. Genetic correlations across geographic regions span US, continental Europe, Nordic, UK/Australia; correlations across age groups span younger (age ≤ 25), midlife (25-64) and older adults (65+).




Figure 1. Manhatt plots of Genome-Wide Associations for each Big Five personality trait (A-E) and genetic correlations across traits (F). Genome wide significance is denoted with a red line at $p = 5 \times 10^{-8}$. A selection of genes containing or nearby the most significant lead SNPs are annotated in each panel (Supplementary Tables S23-S27). Genetic correlations in panel F were estimated among EUR participants using LDSC; standard errors are in parentheses.

Genetic variants with previously reported significant Big Five associations evinced a high replication rate: 207 of 246 (84%) lead SNPs identified in the most recent Big Five GWAS, which partially overlapped with the present GWAS sample, were found in the same locus as a significant variant in this study (Gupta et al., 2024; **Supplementary Table S13**). Individual SNP effects were extremely small, reflecting personality's highly distributed genetic architecture. For example, the median association estimate among extraversion lead SNPs, corrected for the winner's curse (Forde et al., 2023), is .009 SD per effect allele (corresponding to scoring at the 50.35th vs 50th percentile in extraversion). Genetic effects on personality were especially strongly dispersed, as indicated by Linkage Disequilibrium (LD) Fourth Moments Regression (O'Connor et al., 2019). On average, the estimated effective number of independently associated common SNPs for each trait was 16,180 (**Supplementary Table S14**); Extraversion, agreeableness, conscientiousness, and openness to experience were each estimated to be as or more polygenic than 31 of the 32 biobehavioral phenotypes compared in O'Connor and colleagues (2019), including schizophrenia liability and smoking status. We present additional GWAS results, including genetic ancestry-stratified analyses, **Supplementary Figures S1-S10** and **Supplementary Tables S15-S16**.

Characterizing Common-Variant Heritability

Among EUR participants, SNP heritability (h^2_{SNP}) estimated for the Big Five traits using LD Score Regression (LDSC; Bulik-Sullivan et al., 2015) ranged from 4.8% (SE = 0.2%) for agreeableness to 9.3% (SE = 0.3%) for extraversion (**Table 1**; **Supplementary Table S4**). Importantly, these SNP heritability estimates from GWAS meta-analysis incorporate genetic effects that are consistent across contributing cohorts. To allow for variability in genetic effects across cohorts, we conducted a random-effects meta-analysis of cohort-specific h^2_{SNP} estimates, which indicated an average h^2_{SNP} of 8.6% (SE = 0.6%) across traits (ranging from 7.4% for agreeableness to 10.6% for extraversion; **Table 1**; **Supplementary Table S17**), with significant variability across cohorts (mean τ = 3.6%).

h^2_{SNP} is expected to vary as an inverse function of the measurement error in the measurement instrument used (Spearman, 1910; Tucker-Drob, 2017). We confirmed this to be the case using a weighted meta-regression in which we regressed cohort-level h^2_{SNP} on Cronbach's alpha (which indexes a measure's internal consistency on a scale of 0-1), allowing for random intercepts to account for nesting of trait estimates within cohorts. Across all traits, we found that personality measures with greater internal consistency tended to be much more heritable (b = 6.7%, SE = .7%; **Supplementary Figure S11**). In this analysis, the expected h^2_{SNP} for an error-free personality measure ranged from 10.5% for agreeableness (SE = 1.0%) to 16.2% for extraversion (SE = 1.0%; **Table 1**).

Biological follow-up of GWAS signal using MAGMA (de Leeuw et al., 2015) indicated that enriched gene-sets intersected across the Big Five (mean enrichment rank-order ρ = .72; **Supplementary Figures S12-S17**), providing evidence for trait-overlapping molecular and cellular systems in personality neurobiology despite only modest genetic correlations (**Figure 1 panel F**). Consistent with theories of personality development that emphasize the prefrontal cortex (Casey & Caudle, 2013; Romer et al., 2017), genetic associations for each Big Five trait except agreeableness were enriched in genes expressed in the prefrontal cortex (among these, our

top lead SNPs implicate *RCE1*, *FOXP2*, and *SEMA6D*, indicated in **Figure 1; Supplementary Tables S18-S27**). Each of the Big Five demonstrated strong enrichment in protein truncating variant intolerant gene-sets specifically expressed in neurons (e.g., *ARNTL*, *TCF4*, *NEGR1*; **Figure 1**). This pattern, which has also been found for psychiatric disorders and cognitive function (Grotzinger et al., 2022a; Grotzinger et al., 2025), suggests personality-relevant variants are under negative selection, which appears inconsistent with evolutionary theories that posit balancing selection mechanisms as responsible for maintaining genetic variation in personality (Penke & Jokela, 2015). Neurobiological theories of the Big Five posit dopaminergic etiology to variation in openness to experience and extraversion, and serotonergic etiology to variation in conscientiousness, neuroticism, and agreeableness (Carver & Miller, 2006; DeYoung et al., 2021; Depue & Collins, 1999). Based on sets of genes differentially expressed in specific brain cell types, both in mouse and human post-mortem tissues, we found little support for these hypotheses across trait-stratified tests: genes differentially expressed in several types of dopaminergic and serotonergic neurons were in some cases (nominally) significantly enriched for specific outcomes but neither the effect size or p-value placed them among the most enriched neuron types (**Supplementary Tables S18-S22**), suggesting their prominence in the literature relative to other types of neurons is not warranted.

To further characterize the generalizability of genetic effects on personality, we examined the concordance of genetic signal across geography, age, veteran status, measurement instrument, and reporter perspective. We clustered cohorts by these grouping variables, re-estimated GWAS meta-analyses within balanced subgroups, and estimated genetic correlations between subgroups using LDSC (**Table 1; Supplementary Figures S18-S22**). We found that genetic effects were highly similar across four western country clusters (United States, Continental Europe, Nordic, and United Kingdom/Australia, mean $r_g = .86$), three age groups (young [≤ 25], middle [25-64] and older [65+], mean $r_g = .80$), between Million Veteran Program participants and other cohorts (mean $r_g = .80$), and across four personality measurement instruments (mean $r_g = .89$). Additional characterization of genetic architecture across measurement instruments using Genomic Structural Equation Modeling (Grotzinger et al., 2019) confirmed that genetic effects plausibly operate at the level of instrument-general latent factors, with only one locus (associated with both higher extraversion and lower neuroticism) demonstrating significantly heterogeneous effects across measurement instruments (**Supplementary Tables S28-S30, Supplementary Figures S23-S29**). Notably, correlations across geography and measurement instrument were lower for agreeableness than the other Big Five (**Table 1**). This greater heterogeneity explains in part why agreeableness exhibited lower heritability than other traits in the combined meta-analytic GWAS: its genetic effects are less consistent across cohorts. In the Estonian Biobank, where participants' personality was assessed both by their self-report ($N = 73,983$) and reports of close others ($N = 20,269$), we found a high degree of genetic overlap between rater perspectives (mean $r_g = .84$), providing strong evidence that the genetic architecture of personality is not a simple epiphenomenon of self-perception.

Polygenic Prediction of Personality

Past efforts to predict personality from polygenic indices (PGIs) have been hampered by low power of the discovery GWAS used to estimate prediction weights (Becker et al., 2021). Here, we capitalized on the very large sample sizes of our EUR discovery GWAS to estimate weights for PGIs using SBayesR (Lloyd-Jones et al., 2019). PGIs were then tested in five

independent cohorts (holding the cohort out from discovery GWAS when there was sample overlap). In all five cohorts, PGIs predicted significant additive variance in their respective phenotypic personality trait score among EUR participants, after controlling for sex, age, age², and 10 ancestral principal components (extraversion $M_\beta = .19$, range = [.16, .23]; agreeableness $M_\beta = .10$, range = [.07, .14]; conscientiousness $M_\beta = .16$, range = [.12, .18]; neuroticism $M_\beta = .16$, range = [.10, .18]; openness to experience $M_\beta = .17$, range = [.15, .23]; **Figure 2A**). These estimates approximate mean R^2 values ranging from 1.0% for agreeableness to 3.6%, for extraversion. These estimates represent an improvement in prediction over past research (Gupta et al., 2024) for all traits besides agreeableness. Further, highly consistent estimates across cohorts indicates a limited role of birth year, nationality, and ascertainment method in prediction accuracy.

We also predicted personality traits among AFR participants in the Add Health and HRS cohorts, again using weights constructed from the highly powered EUR GWAS. PGI prediction is expected to substantially decrease when there are differences in genetic ancestry between discovery and target samples (Wang et al., 2023). Nevertheless, PGIs significantly predicted extraversion across both cohorts, and neuroticism in HRS ($p < .01$; **Figure 2A**), providing the first evidence for significant PGI prediction of personality traits beyond EUR individuals.




Figure 2. Polygenic prediction and additional genetic associations. Ext = Extraversion. Agr =

Agreeableness. Con = Conscientiousness. Neu = Neuroticism. Ope = Openness to Experience. Panel A depicts standardized prediction of phenotypic personality trait scores among EUR and AFR participants in five cohorts using PGI weights derived from the EUR population-level GWAS. In panel B, betas for continuous traits are standardized and have point estimates depicted with circles, and betas for binary traits are logistic betas and have point estimates depicted with diamonds; percentage affirmative responses are noted in parentheses. † = questions asked only among a subgroup of participants (see Supplementary Text). Panel C depicts correlations between PGIs for each pair of family members. ‡ = Educational Attainment estimates come from Torvik et al. (2022), estimated across spouses in the MoBa cohort. In panel D, accelerometer activity indexes physical movement across a 24-hour period; see Grotzinger et al. (2022b) for complete details. Panel E depicts associations between PGI and census area of residence at age 50+, controlling for birthplace in addition to sex, age, array, and genetic principal components. Only associations that are significant at $p < .01$ after application of Benjamini-Hochberg false-discovery rate correction and sign-concordant within sibship are plotted. In all panels, error bars depict 95% confidence intervals. * = $p < .01$.

Associations with Socially Relevant Behaviors and Important Life Outcomes

Building on epidemiological and longitudinal research that has demonstrated the breadth of personality's associations, we quantified the widespread relevance of personality genetics to a plethora of socially relevant behaviors and important life outcomes (Bleidorn et al., 2019; Roberts et al., 2007; Soto, 2019; Wright et al., 2023) across tests of genetic correlation, PGI prediction, and Mendelian Randomization (MR) applied to EUR GWAS data. Results for AFR GWAS data are reported in the **Supplementary Text** and **Supplementary Figures S30-S31**.

Genetic correlations

Genetic correlations estimated using LDSC indicate widespread genetic sharing between personality traits and health-relevant daily behaviors. Conscientiousness in particular was linked to reduced substance use, greater sports participation, and greater preference for low-calorie foods, as well as the downstream consequences of these behaviors: fewer spells in the hospital, healthier aging, and lower BMI (**Figure 3; Supplementary Tables S31-S32**). Personality was also genetically correlated with fluctuations in accelerometer-measured behavior across the day, with openness to experience linked to increased night-time activity and conscientiousness to increased activity during the day (**Figure 2D; Guerreiro et al., 2024**).



Figure 3. Genetic correlations between the Big Five personality traits and 67 behaviors and outcomes across 9 domains. E = Extraversion. A = Agreeableness. C = Conscientiousness. N = Neuroticism. O = Openness to experience. COVID = Coronavirus disease. HDL = High-density lipoprotein. LDL = Low-density lipoprotein. ADHD = Attention deficit hyperactivity disorder. Letters corresponding to personality traits represent correlation point estimates. Error bars represent 95% confidence intervals. Associations not significant at $p < .01$ are depicted in gray. Associations estimated using LDSC among EUR participants. Full description of each behavior and outcome are available in **Supplementary Tables S31 and S32**.

We found substantial genetic correlations between the Big Five and diagnosis liability for each of 10 psychiatric disorders and 4 transdiagnostic psychiatric disorder factors (**Figure 3**). Genetic profiles associated with neuroticism were associated with increased risk to all forms of psychopathology, particularly internalizing disorders, whereas genetic profiles associated with agreeableness displayed cross-cutting protective associations (Lo et al., 2017). We also identified broad patterns of genetic sharing between openness and psychopathology, which are notably discrepant from the small and inconsistent associations commonly reported in phenotypic research (Kotov et al., 2010; Widiger & Crego, 2019). Many other personality-psychopathology associations were specific and differentiated: Extraversion genetics were associated with lower risk for internalizing disorders but greater risk for neurodevelopmental disorders, and conscientiousness genetics were associated with lower risk for neurodevelopmental disorders but greater risk for compulsive disorders. These findings provide strong support for the close connection between personality traits and psychiatric disorders posited in psychiatric nosologies (Kotov et al., 2017).

Investigation of genetic correlations between personality traits and survey behavior in the UK Biobank indicated that personality plays a major role in research participation (**Figure 3**). In particular, genetic variants associated with openness to experience and agreeableness were associated with completing optional questionnaires, whereas genetic variants associated with neuroticism were associated with fewer questionnaire responses, more “*I don’t know*” and “*prefer no response*” answers, and higher rates of nonresponse after beginning a questionnaire. The genetics of personality may thus have a widespread, underacknowledged influence on sample composition and responses in published research that relies on survey data.

PGI prediction of behaviors and outcomes

We examined PGI prediction of interviewer-rater qualities and delinquency, novel attributes that have not been examined in GWAS, in the Add Health cohort, a representative sample of young US adults ($N = 5,110$). We found that personality PGIs predicted how a young adult was perceived by their in-person interviewer during data collection (**Figure 2B; Supplementary Table S33**): Those with PGIs reflecting lower neuroticism and higher extraversion, agreeableness, and openness to experience were perceived to have a more attractive personality, and those with PGIs reflecting higher agreeableness and conscientiousness were perceived to be more well-groomed. In a subsample of at-risk Add Health participants, personality PGIs were associated with delinquent behavior and intergenerational contact with the prison system, (**Figure 2B; Supplementary Table S33**): high polygenic propensity for neuroticism and low polygenic propensity for agreeableness were associated with school suspension, running away from home, and ever pulling a knife or gun on another person.

Our socio-demographic analysis of UK Biobank participants ($N = 419,261$) linked personality PGIs to the characteristics of one's residence in later adulthood, controlling for place of birth to reflect residential mobility. Focusing only on results that were significant in population-level analyses and sign-concordant in within-family analyses, genetic profiles indicative of higher openness to experience predicted moving into cosmopolitan professional areas and away from suburbia and challenged neighborhoods by middle-older adulthood (age 50+), whereas genetic profiles indicative of higher conscientiousness predicted moving away from cosmopolitan areas and into affluent suburban communities (**Figure 2E, Supplementary Figures S32-S33**). We present additional residential mobility analyses focused on urban/rural migration in the **Supplementary Text** and **Supplementary Figure S34**.

We also quantified personality similarity between spouses over recent generations. To do this, we estimated two PGIs from approximately balanced, non-overlapping halves of the GWAS discovery sample for each Big Five trait. We analyzed these halves with latent variable modelling to estimate genetic correlations among pairs of relatives that correct for PGI estimation error (**Supplementary Figure S36; Supplementary Table S34**). Spousal similarity is expected to induce genetic correlations on assorted traits among offspring that depart from their genetic relatedness, which can bias heritability estimates in standard population-based GWAS (Border et al., 2021; Horwitz et al., 2023). Inferred genetic correlations between parents, estimated using error-corrected PGI similarity meta-analyzed across spouses, sibling pairs, and cousin pairs from UK Biobank ($N_{\text{dyads}} = 177,411$) and deCODE ($N_{\text{dyads}} = 133,358$), indicated minimal historical assortative mating for each of the Big Five traits (Estimated parental $r_g = .01-.04$; **Figure 2C**). Over past generations, birds of a feather have not flocked together: parents were only minimally more genetically similar in their personality traits, on average, than would be expected by chance, and far less genetically similar than has been estimated with respect to educational attainment ($r = .37$; Torvik et al., 2022). We present genetic correlations stratified by relationship and sample in **Supplementary Table S35** and phenotypic parental similarity estimates in the deCODE sample ($N_{\text{pairs}} = 5,317$) in **Supplementary Figure S35**.

Mendelian randomization

To test causality and directionality in associations between personality traits and biobehavioral outcomes, we applied Mendelian Randomization (MR) tests, which leverage SNPs as instrumental variables in a natural experiment (Sanderson et al., 2022). Per MR best practices, we identified a subset of 13 outcomes most likely to comport with core assumptions of this method (**Supplementary Tables S36-S37**), we tested associations across three MR estimators (weighted median, MR-CAUSE, and weighted mode), and we considered associations significant only if they were directionally consistent across all three estimators with 95% confidence/credibility intervals that excluded zero across at least two of the three (see **Supplementary Text** and **Supplementary Figure 38** for complete information on MR methodology).

We found plausible causal effects for 33 exposure-outcome pairings (**Supplementary Table S38**). Twenty-four analyses indicated effects of personality on biobehavioral outcomes, providing novel evidence that personality traits modify physical health – for example, extraversion increased COVID infection risk, conscientiousness decreased BMI and reduced likelihood of smoking initiation and frequency of spells in hospital, and neuroticism decreased

liability for a healthy aging/longevity factor. Furthermore, in line with theoretical perspectives that life experiences affect personality development (Roberts & Yoon, 2022), we found nine plausibly causal effects of biobehavioral outcomes on personality, encompassing broad multi-trait effects of educational attainment, increased BMI, and smoking initiation on personality. Reassuringly, we found no evidence for MR associations between personality and three negative controls (birthweight, number of sisters, and number of brothers) that would only be associated with one's own personality through intergenerational confounding (residual gene-environment correlation). Our highly powered GWAS of each of the Big Five enabled these tests, providing valuable information on the potential causes and consequences of personality traits, which to date have been especially challenging to obtain using other methods (Grosz et al., 2020; Lucas, 2023). Nevertheless, future work should further triangulate on the MR-based causal inferences reported here with additional methods and data that permit strong causal inference (e.g. natural experiments; Bailey et al., 2024; Hammerton & Munafò, 2021).

Evaluating Confounding Caused by Gene-Environment Correlation

Polygenic prediction within dizygotic twin pairs

We used data from the Netherlands Twin Registry ($N_{\text{families}} = 2,956$) and Twins Early Development Study ($N_{\text{families}} = 4,751$) to predict personality traits within dizygotic twin pairs. These models predict personality from twin *differences* in PGIs, which we constructed from the primary EUR population GWAS meta-analysis, excluding the respective twin cohort used for PGI prediction. By holding family environment and parent genotype constant, and leveraging the randomness of intergenerational genetic transmission, these within-family comparisons substantially reduce genetic confounding associated with population stratification and assortative mating that may be present in population-level GWAS (Malanchini et al., 2024; Nivard et al., 2021; Selzam et al., 2019; Veller & Coop, 2024).

For all Big Five traits in both cohorts, within-twin-pair PGI differences robustly predicted the respective personality trait, with indistinguishable magnitude from population-level PGI prediction (Mean within-family $\beta = 100.9\%$ of population-level β , **Figure 4B**). This similarity between methods suggests no detectable confounding in population-level genetic effects on personality. This is in stark contrast with the notable predictive attenuation found in sibling PGI comparisons for other behavioral traits, such as educational attainment (mean within/population predictive strength $\sim 60\%$), cognitive ability ($\sim 80\%$; Okbay et al., 2022; Tan et al., 2024), and externalizing psychopathology ($\sim 75\%$; Karlsson Linnér, 2021).

Direct and indirect genetic effects on personality

We further tested for confounding using a complimentary method. We leveraged parent-offspring data from deCODE ($N = 34,506$) to decompose population-level PGI prediction into direct effects (genetic variants passed down from parent to offspring) and indirect effects (parental genetic variants not transmitted to the offspring). Across the Big Five, direct effects accounted for 96.2% of variance in in population-level PGI prediction, on average (**Figure 4; Supplementary Table S39**), indicating that genetic effects on personality operated nearly exclusive through direct transmission from parent to offspring. Only for conscientiousness were non-transmitted genetic effects significantly predictive of offspring personality (explaining 8.1% of the total predicted effect, $p = .01$). Non-transmitted effects did not differ in magnitude across mothers and fathers (**Supplementary Table S40**). In comparison, for educational attainment,

only 75.1% of variance was explained by direct effects (**Supplementary Table S39**; Kong et al., 2018). After controlling for indirect effects, PGI prediction of extraversion by the extraversion PGI was similar in magnitude to PGI prediction of educational attainment by the educational attainment PGI. Thus, for personality traits, we find a negligible role of parental genetic nurture.




Figure 4. Evaluating environmental confounding in genetic associations with personality. In panel A, the magnitude of PGI prediction from the EUR population-level GWAS is presented for population-level and within-family analyses in two cohorts: Netherlands Twin Registry (NTR) and Twins Early Development Study (TEDS). In panel B, the magnitude of total PGI prediction from the EUR population-level GWAS among offspring in the deCODE cohort (with significance asterisks below the corresponding bar) is decomposed into directly transmitted and indirectly transmitted (non-inherited, with significance asterisks

above the corresponding bar) genetic effects from parents (**Supplementary Table S39**). Comparison estimates for Educational Attainment (EA) come from a PGI constructed from the Okbay et al. (2022) GWAS of EA with deCODE and 23&Me cohorts held out. In panel C, h^2_{SNP} estimates estimated with LDSC from the within-family GWAS (“Wit.”) are presented alongside full EUR population-level GWAS (“Full pop.,” **Table 1**) and matched population-level (“Pop.”) GWAS in cohorts that contributed to within-family GWAS (**Supplementary Table S42**). r_g below each pair of traits indicates their genetic correlation estimated with LDSC, with deviation from $r_g = 1$ tested using a nested chi-square model in Genomic SEM. p -values above each pair of traits indicates results of a test for differences in heritability between population and within-family estimates. In all panels, error bars depict 95% confidence intervals and standard errors are in parentheses. $ns = p \geq .05$. $* = p < .05$. $** = p < .01$. $*** = p < .001$.

Within-family genome-wide analyses

To conduct a maximally stringent test of genetic confounding, we supplemented our population-level GWAS with a within-family GWAS for each Big Five trait. Within-family GWAS accounts for effects of confounding at the level of individual SNPs (Friedman et al., 2021; Tan et al., 2024; Veller & Coop, 2024). To perform within-family GWAS, we assembled data across 12 contributing EUR cohorts ($N = 31,544$ – $50,725$ across traits) that ascertained genetic data among family members using best-practices quality control (Tan et al., 2024), and we combined this with published within-family GWAS estimates of Neuroticism (Howe et al., 2022) (**Supplementary Table S41**; **Supplementary Figures S39–S43**).

For this within-family GWAS, LDSC-estimated h^2_{SNP} estimates ranged from 7.6% (SE = 1.6%) for agreeableness to 13.4% (3.2%) for openness to experience, evincing comparable magnitudes to those reported for the complete meta-analyses of population GWAS (**Figure 4**; **Supplementary Table S42**). To maximize comparability between within-family and population-level GWAS, we also estimated h^2_{SNP} using meta-analysis of population-level GWAS among the matched subset of 12 cohorts who contributed to the within-family GWAS (**Figure 4**). h^2_{SNP} comparisons across models, using $p < .05$ as a significance threshold to ensure identification of potential differences, indicated that for only openness to experience were matched population-level h^2_{SNP} estimates significantly greater than within-family h^2_{SNP} estimates (19.4% vs 13.4%, $p = .02$). This observed difference was driven largely by an inflated population-level h^2_{SNP} estimate for openness to experience relative to the full meta-analysis, and was specifically attributable to the Estonian Biobank sample (**Supplementary Table S41**). In contrast, previously-reported comparisons for other social and behavioral traits, such as educational attainment, depression, and household income, indicate LDSC h^2_{SNP} is attenuated by $\geq 50\%$ in within-family data (Tan et al., 2024). LDSC-estimated genetic correlations between within-family and population-level GWAS summary data averaged $r_g = .86$ and were significantly less than 1.00 only for neuroticism ($r_g = .75$, SE = .08, $p < .001$) and agreeableness ($r_g = .79$, SE = .09, $p = .03$); these results indicate strong correspondence in genetic effect estimates between population-level and within-family analyses. The final piece of evidence for similarity between population-level and within-family genetic effects came from replication analyses of significant and suggestive effects ($p < 1 \times 10^{-5}$) from the full EUR population-level GWAS (excluding matched cohorts): on average, within-family estimates replicated at 95% the magnitude of population-level effect replication (see **Supplementary Table S42** for complete details).

Discussion

In a major consortium effort assembling data from 46 cohorts covering 611K–1.14M EUR and AFR participants, we conducted highly-powered GWAS of each of the Big Five

personality traits, producing dramatic gains in the number of discovered loci, validating powerful and robust polygenic indices, and comprehensively characterizing genetic architecture, confounding, assortative mating, and widespread genetic associations with socially relevant behaviors, health, and important life outcomes. These results overhaul the state of scientific knowledge on the genetic etiology of variation in human personality, establishing a rigorous basis for genetic inference and a fundamental role of personality genetics in the human condition.

Our results reveal that genetic associations with human personality are *generalizable* in several respects. Genetic effects on the Big Five are highly similar across geography, age groups, self-rated versus other-rated report, military service, and measurement instrument; furthermore, PGIs predicted personality traits with near-equivalent magnitude across the lifespan and in samples from different nations. Though the mechanisms by which genetic variance manifests in personality trait differences are sure to be numerous and complex, and may operate in part through dynamic developmental processes (Roberts & Jackson, 2008; Scarr & McCartney 1983), this observed similarity in genetic architecture and PGI prediction across groups suggests that widespread genetic inference is nonetheless possible. One key departure from this overall pattern pertains to agreeableness, which exhibited somewhat lower genetic correlations across geography and rater instrument, suggesting that mechanistic processes relating to this trait may be more varied and inference must be more contextualized.

We also demonstrated that genetic effects on human personality are relatively *unconfounded* by environmental factors that are shared across family members, including uncontrolled population stratification, assortative mating, and dynastic effects, each of which can inflate estimates of genetic effects (Veller & Coop, 2024). PGIs constructed using weights derived from population-level GWAS predicted each of the Big Five traits at indistinguishable strength in a standard population-level analysis and within-DZ twin pair analyses that account for these confounds. Parent-offspring analyses indicated that nearly all prediction was attributable to directly inherited genetic variants, rather than indirect effects of non-transmitted parental genotype. With the exception of openness to experience, estimates of h^2_{SNP} based on population GWAS were indistinguishable from those based on within-family GWAS that accounts for confounding. Effect size replication for top independent loci was very similar among population-level versus within-family hold-out samples. Finally, our comparisons of genetic correlations across relatives indicate that there has been minimal assortative mating on personality traits across recent generational time, extending decades of phenotypic research on romantic partner similarity (e.g. Horwitz et al., 2022; McCrae et al., 2008). Research in social science genomics has indicated that environmental sources of confounding play a major role in population-level GWAS of traits such as educational attainment, income, and cognitive performance (Howe et al., 2022; Okbay et al., 2022; Tan et al., 2024), introducing considerable challenge in obtaining estimates of direct genetic effects for these traits. In contrast, for personality, the lack of confounding indicates that population-level GWAS estimates overwhelmingly reflect direct genetic effects.

Functional genomic analyses and biological annotation indicated that despite their only modest genetic intercorrelations, genetic effects on each of the Big Five occur via overlapping molecular and cellular systems; for example, the prefrontal cortex was implicated in the genetics

of each trait aside from agreeableness. Analyses also provide little support for neurobiological theories of the Big Five that posit trait-differentiated serotonergic and dopaminergic etiologies (Carver & Miller, 2006; DeYoung et al., 2021; Depue & Collins, 1999). Single cell gene expression data integrated with our GWAS suggests there is no straightforward reductive mapping from broad, multifaceted personality traits to specific neuron types. However, the availability of these high powered GWAS combined with ever-expanding access to temporal and spatial brain gene expression data will enable future developmental and system-specific analysis of neural personality etiology. Each of the Big Five demonstrated strong enrichment in protein truncating variant intolerant gene-sets specifically expressed in brain cells, suggesting that genetic variants with strong effects on personality are selected against (i.e. undergo negative selection), which is at odds with evolutionary theories that posit genetic variation in personality is solely maintained by inconsistent and varying selection pressures on personality that are in aggregate directionally neutral (balancing selection) (Penke & Jokela, 2016). Future research will be needed to further clarify the relative contributions of negative and balancing selection to variation in human personality traits.

Key limitations point to focal areas for future research. First, our within-family GWAS meta-analysis, while the largest of personality to date, was small compared to the population GWAS meta-analysis. Collection of additional within-family data, especially intergenerational family units (Davies et al., 2024) will provide further leverage to accurately and precisely identify loci indexing direct genetic effects. Fine mapping and experimental designs (e.g., Sanchez-Roige et al., 2023) will also be necessary to identify the causal variants within the identified loci. Second, we examined genetic correlations across only broad stratifying variables, and only among EUR and AFR individuals. More fine-grained sex- and age- stratified analyses and stratification by more expansive ancestral and cultural backgrounds will further refine inferences about the generalizability and differentiation of genetic effects on personality. Finally, we focused only on the very broad Big Five personality traits. Research with more granular personality trait data (e.g., of specific behavioral tendencies; Möttus et al., 2017) will provide greater specificity of inference and permit more sophisticated multivariate analysis of the genetic patterning of personality trait structure.

Much of the scientific value of studying personality stems from its applicability to nearly all aspects of the lives that people lead. We extend this tenet by linking heritable variation in personality traits to core mental and physical health outcomes, labor market performance, and reproduction, with differentiated patterns that implicate certain traits (e.g. conscientiousness) as especially relevant to certain classes of outcomes (e.g. health behaviors). The genetics of personality also predict patterns of everyday behavior: how and when a person responds to surveys, where they move throughout their lives, and even the impression they make on others in a single interview session. Personality traits are highly stable over time (Bleidorn et al., 2022) and are resistant to simple experimental manipulation (Briley et al., 2018). Causal inference with respect to personality's associations afforded by robust MR analyses is therefore especially valuable. MR results implicate personality traits as both causes and consequences of a range of health conditions. Given the pervasive relevance of personality to human experience, researchers across scientific fields will benefit from incorporating personality traits into models and theories when predicting and explaining biopsychosocial outcomes.

Acknowledgements

T.S., M.G.N., and E.M.T-D. were supported by NIMH R01MH120219. E.M.T-D. was also supported by NIA R01AG073593. W.D.H., and C.X. were supported by a Career Development Award from the Medical Research Council (MRC) [MR/T030852/1] for the project titled “From genetic sequence to phenotypic consequence: genetic and environmental links between cognitive ability, socioeconomic position, and health.” T.K. was funded by the European Research Council (ERC Consolidator Grant awarded under the Horizon Europe framework, grant agreement 101087395. F.S. was supported by the Hector foundation II and by a 2023 NARSAD Young Investigator Grant (#31537) from the Brain & Behavior Research Foundation with support from the Families for Borderline Personality Disorder Research. R.C. was supported by the Jacobs Foundation grant no. 2023-1510-00 and the Research Council of Norway (grant number 325245). H.A. was supported by the research council of Norway (#274611, #324620). F.A.T. was supported by Research Council of Norway #26270. E.C.C. was supported by the research council of Norway (RCN; #274611) and the South-Eastern Norway Regional Health Authority (HSØ; #2021045). J.L. received financial support from the Strategic Research Council (SRC) established within the Academy of Finland (decision number: 352700). K.I. and U.Va. have been funded by Estonian Research Council's personal research funding start-up grants PSG656 and PSG759. R.M. has been funded by Estonian Research Council's team grant PRG2190. U.Võ. and T.E. have been funded by Estonian Research Council's team grant PRG1291.

References

- Anni, K., Vainik, U., & Möttus, R. (2024). Personality profiles of 263 occupations. *Journal of Applied psychology*.
- Antonoplis, S., & John, O. P. (2022). Who has different-race friends, and does it depend on context? Openness (to other), but not agreeableness, predicts lower racial homophily in friendship networks. *Journal of personality and social psychology*, 122(5), 894-919.
- Bailey, D. H., Jung, A. J., Beltz, A. M., Eronen, M. I., Gische, C., Hamaker, E. L., ... & Murayama, K. (2024). Causal inference on human behaviour. *Nature human behaviour*, 8(8), 1448-1459.
- Beck, E. D., & Jackson, J. J. (2022). A mega-analysis of personality prediction: Robustness and boundary conditions. *Journal of personality and social psychology*, 122(3), 523-553.
- Becker, J., Burik, C. A., Goldman, G., Wang, N., Jayashankar, H., Bennett, M., ... & Okbay, A. (2021). Resource profile and user guide of the Polygenic Index Repository. *Nature human behaviour*, 5(12), 1744-1758.
- Bleidorn, W., Hill, P. L., Back, M. D., Denissen, J. J., Hennecke, M., Hopwood, C. J., ... & Roberts, B. (2019). The policy relevance of personality traits. *American psychologist*, 74(9), 1056-1067.
- Bleidorn, W., Schwaba, T., Zheng, A., Hopwood, C. J., Sosa, S. S., Roberts, B. W., & Briley, D. A. (2022). Personality stability and change: A meta-analysis of longitudinal studies. *Psychological bulletin*, 148(7-8), 588-619.
- Bogg, T., & Roberts, B. W. (2004). Conscientiousness and health-related behaviors: a meta-analysis of the leading behavioral contributors to mortality. *Psychological bulletin*, 130(6), 887-919.
- Borghans, L., Golsteyn, B. H., Heckman, J. J., & Humphries, J. E. (2016). What grades and achievement tests measure. *Proceedings of the National Academy of Sciences*, 113(47), 13354-13359.
- Briley, D. A., Livengood, J., & Derringer, J. (2018). Behaviour genetic frameworks of causal reasoning for personality psychology. *European Journal of Personality*, 32(3), 202-220.
- Bucher, M. A., Suzuki, T., & Samuel, D. B. (2019). A meta-analytic review of personality traits and their associations with mental health treatment outcomes. *Clinical psychology review*, 70, 51-63.
- Bulik-Sullivan, B. K., Loh, P. R., Finucane, H. K., Ripke, S., Yang, J., Schizophrenia Working Group of the Psychiatric Genomics Consortium, ... & Neale, B. M. (2015). LD Score regression distinguishes confounding from polygenicity in genome-wide association studies. *Nature genetics*, 47(3), 291-295.
- Carver, C. S., & Miller, C. J. (2006). Relations of serotonin function to personality: Current views and a key methodological issue. *Psychiatry research*, 144(1), 1-15.
- Casey, B. J., & Caudle, K. (2013). The teenage brain: Self control. *Current directions in psychological science*, 22(2), 82-87.
- De Leeuw, C. A., Mooij, J. M., Heskes, T., & Posthuma, D. (2015). MAGMA: generalized gene-set analysis of GWAS data. *PLoS computational biology*, 11(4), e1004219.
- De Moor, M. H., Beem, A. L., Stubbe, J. H., Boomsma, D. I., & De Geus, E. J. (2006). Regular exercise, anxiety, depression and personality: a population-based study. *Preventive medicine*, 42(4), 273-279.
- Deary, I. J., Weiss, A., & Batty, G. D. (2010). Intelligence and personality as predictors of illness and death: How researchers in differential psychology and chronic disease epidemiology are collaborating to understand and address health inequalities. *Psychological science in the public interest*, 11(2), 53-79.
- Depue, R. A., & Collins, P. F. (1999). Neurobiology of the structure of personality: Dopamine, facilitation of incentive motivation, and extraversion. *Behavioral and brain sciences*, 22(3), 491-517.
- DeYoung, C. G., Grazioplene, R. G., & Allen, T. A. (2021). The neurobiology of personality. In O. P. John & R. W. Robins (Eds.), *Handbook of personality: Theory and research* (pp. 193-216). The Guilford Press.
- Duckitt, J., & Sibley, C. G. (2010). Personality, ideology, prejudice, and politics: A dual-process motivational model. *Journal of personality*, 78(6), 1861-1894.
- Ebert, T., Gebauer, J. E., Brenner, T., Bleidorn, W., Gosling, S. D., Potter, J., & Rentfrow, P. J. (2022). Are regional differences in psychological characteristics and their correlates robust? Applying spatial-analysis techniques to examine regional variation in personality. *Perspectives on psychological science*, 17(2), 407-441.
- Forde, A., Hemani, G., & Ferguson, J. (2023). Review and further developments in statistical corrections for Winner's Curse in genetic association studies. *PLoS genetics*, 19(9), e1010546.
- Friedman, N. P., Banich, M. T., & Keller, M. C. (2021). Twin studies to GWAS: there and back again. *Trends in Cognitive sciences*, 25(10), 855-869.
- Guerreiro, J., Schulze, L., Garcia i Tormo, A., Henwood, A. J., Schneider, L., Krob, E., ... & Matic, A. (2024). The Relationship Between Big Five Personality Traits and Sleep Patterns: A Systematic Review. *Nature and science of sleep*, 1327-1337.

- Graham, E. K., Rutsohn, J. P., Turiano, N. A., Bendayan, R., Batterham, P. J., Gerstorf, D., ... & Mroczek, D. K. (2017). Personality predicts mortality risk: An integrative data analysis of 15 international longitudinal studies. *Journal of research in personality*, 70, 174-186.
- Grotzinger, A. D., de la Fuente, J., Davies, G., Nivard, M. G., & Tucker-Drob, E. M. (2022a). Transcriptome-wide and stratified genomic structural equation modeling identify neurobiological pathways shared across diverse cognitive traits. *Nature Communications*, 13, 6280.
- Grotzinger, A. D., Mallard, T. T., Akingbuwa, W. A., Ip, H. F., Adams, M. J., Lewis, C. M., ... & Nivard, M. G. (2022b). Genetic architecture of 11 major psychiatric disorders at biobehavioral, functional genomic and molecular genetic levels of analysis. *Nature Genetics*, 54(5), 548-559.
- Grotzinger, A. D., Werme, J., Peyrot, W. J., Frei, O., de Leeuw, C., Bicks, L. K., ... & Smoller, J. W. (2025). The Landscape of Shared and Divergent Genetic Influences across 14 Psychiatric Disorders. *medRxiv*, 2025-01.
- Grotzinger, A. D., Rhemtulla, M., de Vlaming, R., Ritchie, S. J., Mallard, T. T., Hill, W. D., ... & Tucker-Drob, E. M. (2019). Genomic structural equation modelling provides insights into the multivariate genetic architecture of complex traits. *Nature human behaviour*, 3(5), 513-525.
- Gupta, P., Galimberti, M., Liu, Y., Beck, S., Wingo, A., Wingo, T., ... & Levey, D. F. (2024). A genome-wide investigation into the underlying genetic architecture of personality traits and overlap with psychopathology. *Nature human behaviour*, 8, 2235-2249.
- Hammerton, G., & Munafò, M. R. (2021). Causal inference with observational data: the need for triangulation of evidence. *Psychological medicine*, 51(4), 563-578.
- Hampson, S. E., Goldberg, L. R., Vogt, T. M., & Dubanoski, J. P. (2007). Mechanisms by which childhood personality traits influence adult health status: educational attainment and healthy behaviors. *Health psychology*, 26(1), 121-125.
- Hampson, S. E. (2012). Personality processes: Mechanisms by which personality traits “get outside the skin”. *Annual review of psychology*, 63(1), 315-339.
- Horwitz, T. B., Balbona, J. V., Paulich, K. N., & Keller, M. C. (2023). Evidence of correlations between human partners based on systematic reviews and meta-analyses of 22 traits and UK Biobank analysis of 133 traits. *Nature human behaviour*, 7(9), 1568-1583.
- Howe, L. J., Nivard, M. G., Morris, T. T., Hansen, A. F., Rasheed, H., Cho, Y., ... & Davies, N. M. (2022). Within-sibship genome-wide association analyses decrease bias in estimates of direct genetic effects. *Nature genetics*, 54(5), 581-592.
- John, O. P., Naumann, L. P., & Soto, C. J. (2008). Paradigm shift to the integrative big five trait taxonomy. *Handbook of Personality: Theory and Research*, 3(2), 114-158.
- Jokela, M. (2020). Selective residential mobility and social influence in the emergence of neighborhood personality differences: Longitudinal data from Australia. *Journal of research in personality*, 86, 103953.
- Jokela, M., Bleidorn, W., Lamb, M. E., Gosling, S. D., & Rentfrow, P. J. (2015). Geographically varying associations between personality and life satisfaction in the London metropolitan area. *Proceedings of the National Academy of Sciences*, 112(3), 725-730.
- Karlsson Linnér, R., Mallard, T. T., Barr, P. B., Sanchez-Roige, S., Madole, J. W., Driver, M. N., ... & Dick, D. M. (2021). Multivariate analysis of 1.5 million people identifies genetic associations with traits related to self-regulation and addiction. *Nature neuroscience*, 24(10), 1367-1376.
- Kotov, R., Gamez, W., Schmidt, F., & Watson, D. (2010). Linking “big” personality traits to anxiety, depressive, and substance use disorders: a meta-analysis. *Psychological bulletin*, 136(5), 768-821.
- Kotov, R., Krueger, R. F., Watson, D., Achenbach, T. M., Althoff, R. R., Bagby, R. M., ... & Zimmerman, M. (2017). The Hierarchical Taxonomy of Psychopathology (HiTOP): A dimensional alternative to traditional nosologies. *Journal of abnormal psychology*, 126(4), 454-477.
- Landis, B. (2016). Personality and social networks in organizations: A review and future directions. *Journal of organizational behavior*, 37, S107-S121.
- Lloyd-Jones, L. R., Zeng, J., Sidorenko, J., Yengo, L., Moser, G., Kemper, K. E., ... & Visscher, P. M. (2019). Improved polygenic prediction by Bayesian multiple regression on summary statistics. *Nature communications*, 10(1), 5086.
- Lo, M. T., Hinds, D. A., Tung, J. Y., Franz, C., Fan, C. C., Wang, Y., ... & Chen, C. H. (2017). Genome-wide analyses for personality traits identify six genomic loci and show correlations with psychiatric disorders. *Nature genetics*, 49(1), 152-156.

- Malanchini, M., Allegrini, A. G., Nivard, M. G., Biroli, P., Rimfeld, K., Cheesman, R., ... & Plomin, R. (2024). Genetic associations between non-cognitive skills and academic achievement over development. *Nature human behaviour*, 1-13.
- Matthews, G., Deary, I. J., & Whiteman, M. C. (2009). *Personality traits*. Cambridge University Press.
- McCrae, R. R., Martin, T. A., Hrebicková, M., Urbánek, T., Boomsma, D. I., Willemsen, G., & Costa Jr, P. T. (2008). Personality trait similarity between spouses in four cultures. *Journal of personality*, 76(5), 1137-1164.
- Möttus, R., Kandler, C., Bleidorn, W., Riemann, R., & McCrae, R. R. (2017). Personality traits below facets: The consensual validity, longitudinal stability, heritability, and utility of personality nuances. *Journal of Personality and Social Psychology*, 112(3), 474-490.
- National Academies of Sciences, Engineering, and Medicine. (2023). *Using population descriptors in genetics and genomics research: A new framework for an evolving field*. Washington, DC: The National Academies Press. <https://doi.org/10.17226/26902>.
- Nivard, M. G., Belsky, D. W., Harden, K. P., Baier, T., Andreassen, O. A., Yström, E., ... & Lyngstad, T. H. (2024). More than nature and nurture, indirect genetic effects on children's academic achievement are consequences of dynastic social processes. *Nature human behaviour*, 8(4), 771-778.
- O'Connor, L. J., Schoech, A. P., Hormozdiari, F., Gazal, S., Patterson, N., & Price, A. L. (2019). Extreme polygenicity of complex traits is explained by negative selection. *The American journal of human genetics*, 105(3), 456-476.
- Okbay, A., Wu, Y., Wang, N., Jayashankar, H., Bennett, M., Nehzati, S. M., ... & Young, A. I. (2022). Polygenic prediction of educational attainment within and between families from genome-wide association analyses in 3 million individuals. *Nature genetics*, 54(4), 437-449.
- Penke, L., & Jokela, M. (2016). The evolutionary genetics of personality revisited. *Current opinion in psychology*, 7, 104-109.
- Roberts, B. W., Kuncel, N. R., Shiner, R., Caspi, A., & Goldberg, L. R. (2007). The power of personality: The comparative validity of personality traits, socioeconomic status, and cognitive ability for predicting important life outcomes. *Perspectives on psychological science*, 2(4), 313-345.
- Roberts, B. W., & Yoon, H. J. (2022). Personality psychology. *Annual review of psychology*, 73(1), 489-516.
- Romer, D., Reyna, V. F., & Satterthwaite, T. D. (2017). Beyond stereotypes of adolescent risk taking: Placing the adolescent brain in developmental context. *Developmental cognitive neuroscience*, 27, 19-34.
- Sanchez-Roige, S., Jennings, M. V., Thorpe, H. H., Mallari, J. E., van der Werf, L. C., Bianchi, S. B., ... & Palmer, A. A. (2023). CADM2 is implicated in impulsive personality and numerous other traits by genome-and phenome-wide association studies in humans and mice. *Translational Psychiatry*, 13(1), 167.
- Sanderson, E., Glymour, M. M., Holmes, M. V., Kang, H., Morrison, J., Munafò, M. R., ... & Davey Smith, G. (2022). Mendelian randomization. *Nature reviews methods primers*, 2(1), 6.
- Scarr, S., & McCartney, K. (1983). How people make their own environments: A theory of genotype→ environment effects. *Child development*, 424-435.
- Selzam, S., Ritchie, S. J., Pingault, J. B., Reynolds, C. A., O'Reilly, P. F., & Plomin, R. (2019). Comparing within- and between-family polygenic score prediction. *The American journal of human genetics*, 105(2), 351-363.
- Spearman, C. (1910). Correlation calculated from faulty data. *British journal of psychology*, 3(3), 271.
- Soto, C. J. (2019). How replicable are links between personality traits and consequential life outcomes? The life outcomes of personality replication project. *Psychological science*, 30(5), 711-727.
- Soto, C. J., & John, O. P. (2017). The next Big Five Inventory (BFI-2): Developing and assessing a hierarchical model with 15 facets to enhance bandwidth, fidelity, and predictive power. *Journal of personality and social psychology*, 113(1), 117-143.
- Tan, T., Jayashankar, H., Guan, J., Nehzati, S. M., Mir, M., Bennett, M., ... & Young, A. S. (2024). Family-GWAS reveals effects of environment and mating on genetic associations. *medRxiv*, 2024-10.
- Torvik, F. A., Eilertsen, E. M., Hannigan, L. J., Cheesman, R., Howe, L. J., Magnus, P., ... & Ystrom, E. (2022). Modeling assortative mating and genetic similarities between partners, siblings, and in-laws. *Nature communications*, 13(1), 1108.
- Tucker-Drob, E. M. (2017). Measurement error correction of genome-wide polygenic scores in prediction samples. *BioRxiv*, 165472.
- Veller, C., & Coop, G. M. (2024). Interpreting population- and family-based genome-wide association studies in the presence of confounding. *PLoS biology*, 22(4), e3002511.
- Wang, Y., Kanai, M., Tan, T., Kamariza, M., Tsuo, K., Yuan, K., ... & Martin, A. R. (2023). Polygenic prediction across populations is influenced by ancestry, genetic architecture, and methodology. *Cell genomics*, 3(10).

- Widiger, T. A., & Crego, C. (2019). HiTOP thought disorder, DSM-5 psychoticism, and five factor model openness. *Journal of research in personality*, 80, 72-77.
- Widiger, T. A., Sellbom, M., Chmielewski, M., Clark, L. A., DeYoung, C. G., Kotov, R., ... & Wright, A. G. (2019). Personality in a hierarchical model of psychopathology. *Clinical psychological science*, 7(1), 77-92.
- Willroth, E. C., Luo, J., Atherton, O. E., Weston, S. J., Drewelies, J., Batterham, P. J., ... & Graham, E. K. (2023). Personality traits and health care use: A coordinated analysis of 15 international samples. *Journal of personality and social psychology*, 125(3), 629.
- Wright, A. J., & Jackson, J. J. (2023). Do changes in personality predict life outcomes?. *Journal of personality and social psychology*. 125(6), 1495–1518.
- Yoneda, T., Lozinski, T., Turiano, N., Booth, T., Graham, E. K., Mroczek, D., & Terrera, G. M. (2023). The Big Five personality traits and allostatic load in middle to older adulthood: A systematic review and meta-analysis. *Neuroscience & biobehavioral reviews*, 148, 105145.
- Young, A. I., Nehzati, S. M., Benonisdottir, S., Okbay, A., Jayashankar, H., Lee, C., ... & Kong, A. (2022). Mendelian imputation of parental genotypes improves estimates of direct genetic effects. *Nature genetics*, 54(6), 897-905.